

Studying urban land-atmospheric interactions by coupling an urban canopy model with a single column atmospheric model



Central Arizona-Phoenix
Long-Term Ecological Research
CAP LTER


Jiyun Song, Jiachuan Yang and Zhihua Wang
School of Sustainable Engineering and the Built Environment, Arizona State University,
PO Box 875306, Tempe, AZ 85287-5306 (Email: jsong52@asu.edu; zhwang@asu.edu)

Problem Statement


Land use land cover changes (LULCC) in urban area will modify surface energy budgets and turbulent fluxes, which will further change the dynamic and thermodynamic structures of the overlying atmospheric boundary layer (ABL).

Question: How will the overlying atmosphere be influenced by urban LULCC?

Coupling Methodology




Single Column Model (SCM)




2005/03/26 9:30am Point Reyes, CA

SCM-Slab Model Coupling



Wangara Day 33 12:00 pm 3:00pm in Australia


SCM-SLUCM Coupling



Temporal evolution:
The height of atmospheric boundary layer (ABL) will increase due to the influence of turbulent heat fluxes at daytime. The virtual potential temperature, specific humidity, and wind speed will change due to effects of surface fluxes at daytime.


Spatial evolution:
The values of virtual potential temperature and specific humidity are almost constant in the mixed layer.

Effects of geometry




Higher building (shadow effects, incident angle impacts) will lead to:
➢ higher ABL
➢ higher virtual potential temperature
➢ slightly lower specific humidity

Effects of vegetation fraction




Higher vegetation fraction will lead to:
➢ lower ABL
➢ lower virtual potential temperature
➢ higher specific humidity.

Effects of roughness length of momentum



Higher roughness length of momentum above roof will lead to:
➢ higher ABL
➢ higher virtual potential temperature
➢ higher specific humidity.

Effects of roof albedo



Higher roof albedo will lead to:
➢ lower ABL
➢ lower virtual potential temperature
➢ slightly higher specific humidity.

2012/06/13 in pre-monsoon season at Phoenix, AZ

Conclusions and Perspectives

- ❖ We implemented an advanced urban canopy model (UCM) with an improved representation of urban hydrological processes.
- ❖ We simulated urban land – atmosphere interactions for different urban land surfaces with different geometry, vegetation fraction, roughness of length for momentum, and albedo.
- ❖ We are implementing our advanced UCM into WRF to improve regional climate modeling.

Acknowledgements

The authors would like to acknowledge the Central Arizona-Phoenix Long-Term Ecological Research (CAP-LTER) for providing Graduate Grants (CAP3: BCS-1026865) and Dr. Winston Chow for providing the data of CAP-LTER flux tower in Phoenix, AZ.

References

- [1] Z.H. Wang, E. Bou-Zeid, A.S. James (2013). A coupled energy transport and hydrological model for urban canopies evaluated using a wireless sensor network. *Quart J Roy Meteor Soc*, 139 (675): 1643-1657.
- [2] Y. Noh, W. G. Cheon, and S. Y. Hong (2003). Improvement of the K-profile model for the planetary boundary layer based on large eddy simulation data. *Boundary-Layer Meteorol*, 107: 401-427.
- [3] S.Y. Hong, Y. Noh, and J. Dudhia (2006). A new vertical diffusion package with an explicit treatment of entrainment processes. *Mon Weather Rev*, 134: 2318-2341.
- [4] A. J. Dyer (1974). A review of flux-profile relationships. *Boundary-Layer Meteorol*, 7: 363-372.