


The Effect of Urban Heat and Food Availability on Black Widow Spiders

Sarah E. Lindley, Natalija Bogdanovic, Helena Hanna, & J. Chadwick Johnson

School of Mathematical & Natural Sciences, Arizona State University, West



Introduction

- Urbanization creates unique ecosystems, and can alter species diversity and organismal life history [1].
- The Urban Heat Island (UHI) effect results from using heat retentive building materials and causes elevated temperatures [2].
- Urbanized deserts often have a higher resource and prey availability than desert areas [3].
- Black widow spiders (*Latrodectus hesperus*) are a medically-important species [4] that often infest urban habitats [5].
- Urban black widow microclimates are 6°C hotter than desert areas. In the lab, this UHI hurts longevity, growth, and mass [6].
- We predicted that combining urban prey availability and elevated UHI temperatures would favor longevity, growth, and mass.

Methods

- Adult female spiders were collected from urban habitat across Maricopa county.
- Egg sacs were collected from 16 of these spiders, and 100 eggs from each egg sac were weighed (μg) and imaged (mm^2).
- Individual eggs were placed in transparent boxes (4x4x5cm), and reared at 24°C until hatched.
- At day 30, hatched spiderlings from each family were divided between temperature treatments (27°C =desert; 33°C=UHI).
- Half of the spiderlings from each temperature treatment were fed 8 flies per week (low food), and half were fed 16 flies per week (high food).
- Survival, development speed, mass at day 105, and the sex of each spiderling were included as measures in our analyses.

Results

- Family had a significant effect on both longevity and developmental speed (see Figure 1).
- Food and temperature each independently affected developmental speed, but we found no interaction between these variables (see Figure 2).
- Food and temperature each independently affected body mass, but we found no interaction between these variables (see Figure 3).
- Food had a significant effect on longevity. A high food diet resulted in an increase in longevity of an average of 2.375 days (see Figure 4).
- The sex of a spiderling did not have a significant effect on any of the variables analyzed.

Discussion

- UHI temperatures and low food each hurt development, mass, & longevity.
- However, we did not find the predicted interaction where UHI conditions favor black widows when food is abundant.
 - We still have no explanation for why UHI conditions in the laboratory limit black widows that clearly thrive under urban field conditions.
- Perhaps the spatial and/or temporal complexity of urban habitat provides a buffer from extreme temperatures not seen from laboratory protocols.
- Family of origin continues to be an important variable in predicting the black widow's life history.
 - Despite manipulation of both temperature and food availability, variation within sibships continues to be smaller than between sibships.




Fig. 1 Family Affects Development




Fig. 2 Food & Temperature Affect Development




Fig. 3 Food and Temperature Affect Mass




Figure 4. Food Affects Longevity

Literature Cited

- Chick LD, Strickler SA, Perez A, Martin RA, Diamond SE. (2019) Urban heat islands advance the timing of reproduction in a social insect. *Journal of Thermal Biology*, 80: 119-125. <https://doi.org/10.1016/j.jtherbio.2019.01.004>.
- Chow, W. T. L., Brennan, D., Brazel, A. J., Chow, W. T. L., Brennan, D., & Brazel, A. J. (2012). Urban Heat Island Research in Phoenix, Arizona: Theoretical Contributions and Policy Applications. *Bulletin of the American Meteorological Society*, 93(4), 517-530. <https://doi.org/10.1175/BAMS-D-11-00011.1>
- Johnson, J., Trubl, P., & Miles, L. (2012). Black Widows in an Urban Desert: City-Living Compromises Spider Fecundity and Egg Investment Despite Urban Prey Abundance. *The American Midland Naturalist*, 168(2), 333-340. Retrieved from www.jstor.org/stable/23269823
- Vetter, R., & Visscher, P. (1998). Bites and stings of medically important venomous arthropods. *International Journal of Dermatology*, 37(7), 481-496.
- Trubl, P., Gburek, T., Miles, L., & Johnson, J. (2012). Black widow spiders in an urban desert: Population variation in an arthropod pest across metropolitan Phoenix, AZ. *Urban Ecosystems*, 15(3), 599-609.
- Johnson, J. C., Urcuyo, J., Moen, C., & Stevens, D. R., 2nd (2019). Urban heat island conditions experienced by the Western black widow spider (*Latrodectus hesperus*): Extreme heat slows development but results in behavioral accommodations. *PLoS one*, 14(9), e0220153. doi:10.1371/journal.pone.0220153

Acknowledgements

- Special thanks to the members of Dr. Johnson's black widow research lab who helped rear the spiders used in this study.
- Additional thanks to REU CAP LTER for providing funding to help facilitate the accomplishment of this research.

